

Věra Šedivá

On convergence of sequences of functions

Commentationes Mathematicae Universitatis Carolinae, Vol. 2 (1961), No. 3, 3–12

Persistent URL: <http://dml.cz/dmlcz/104889>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1961

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON CONVERGENCE OF SEQUENCES OF FUNCTIONS

V. ŠEDIVÁ - TRNKOVÁ, Praha

If (Q_λ, u_λ) ($\lambda \in \Lambda$) are spaces, we can define on a cartesian product Q of sets Q_λ a convergence of sequences by a well known way: For $x^n, x \in Q$ ($n = 1, 2, \dots$), $x^n \xrightarrow{u} x$ if and only if $x_\lambda^n \xrightarrow{u_\lambda} x_\lambda$ in the space (Q_λ, u_λ) for every $\lambda \in \Lambda$ (x_λ denotes the λ -th coordinate of the point x). This convergence defines a topology u on Q in the well known way (for $A \subset Q$ $u A$ consists of all $x \in Q$ such that $x^n \xrightarrow{u} x$ for some $x^n \in A$). Following J. Novák [3], we call (Q, u) an \mathcal{L} -product of spaces (Q_λ, u_λ) and denote $(Q, u) = \mathcal{L}_{\lambda \in \Lambda} (Q_\lambda, u_\lambda)$. Let us point out that, following E. Čech [1] a topology u on the set Q is defined as a mapping u , which to every $M \subset Q$ assigns a set $u M \subset Q$ and satisfies the following axioms: $u \emptyset = \emptyset$, $u(x) = (x)$, $u(M_1 \cup M_2) = u M_1 \cup u M_2$. The condition $u(u M) = u M$, called axiom F by E. Čech, is not required in general; if it is satisfied, then u is called an F -topology and (Q, u) an F -space; if it does not hold, then u is called a non- F -topology and (Q, u) a non- F -space.

For any topology u on Q two further F -topologies are defined: \tilde{u} , the F -reduction of u , which has an open base consisting of all $Q - u A$, $A \subset Q$; u^* , the F -modification of u , which is the finest of all F -topologies, coarser than u . Clearly $\tilde{u} = u$ or $u = u^*$ if and only if u is an F -topology.

In this note an \mathcal{L} -product of two-point spaces is studied. The smallest cardinal number \aleph_1 is found, for which an \mathcal{L} -product of \aleph_1 two-point spaces is non- F -space, event. it is not countably compact.

It is shown, that the \mathcal{L} -product (Q, u) of uncountable number of two-point spaces and (Q, u^*) are not regular. Several criteria are given, when the space (Q, \tilde{u}) is discrete (I. - III.).

In IV.- VIII. similar questions for subspaces of the space of real-valued functions on some F -space are studied.

In the whole note proofs are omitted.

In this note, N denotes the set of all natural numbers. If A, B are sets, A^B denotes the set of all mappings of B into A . If $\alpha \in A^B$, $x \in B$ then the element, corresponding to x in the mapping α , is denoted $\alpha(x)$ or α_x .

If $\alpha \in A^B$, $C \subset B$, then $\alpha|C$ denotes the mapping of C into A , for which $\alpha|C(x) = \alpha(x)$ for all $x \in C$. \aleph denotes an arbitrary cardinal number.

I. Countable compactness.

Definition:

The space (Q, u) is called countably compact, if every infinite subset of Q has a cluster point.

Theorem 1, 1:

Let (Q, u) be a space. The following properties are equivalent:

- (1) (Q, u) is countably compact.
- (2) (Q, u^*) is countably compact.

Theorem 1,1 does not hold for compactness only.

Definition.

Let σ be the smallest power of a system \mathcal{A} of subsets of N , which has the following property:

if $S \subset N$ is infinite, then there exists $A \in \mathcal{A}$ such that the sets $S \cap A$, $S - A$ are infinite.

Theorem 1, 2:

Let (Q, u) be an \mathcal{L} -product of \aleph two-point spaces.

The following properties are equivalent:

- (1) (Q, u) is not countably compact.
- (2) $\aleph \geq \sigma$.

II. F -axiom, order and regularity.

Definitions:

Let A be an infinite countable set, $\alpha, \beta \in N^A$. We write $\alpha \succ \beta$ if $\alpha(x) > \beta(x)$ for all $x \in A$, except a finite number. If $\alpha \succ \beta$ does not hold, we write $\alpha \not\succ \beta$.

We say that $\mathcal{A} \subset N^A$ is an unbounded system in N^A if for every $\gamma \in N^A$ there exists some $\alpha \in \mathcal{A}$ such that $\gamma \not\succ \alpha$.

We say that $\mathcal{A} = \{\alpha^\lambda\} \subset N^N$ is a hereditary unbounded system, if the system $\{\alpha^\lambda \mid A\}$ is unbounded in N^A for every infinite $A \subset N$.

We say that a set $\mathcal{A} \subset N^N$ is a chain, if it is linearly ordered by the relation \succ .

An unbounded system, which is also a chain, is called an unbounded chain. The existence of an unbounded chain follows from Zorn's lemma.

Definition:

Let τ_1 be the smallest power of unbounded chain.

Let τ_2 be the smallest power of hereditary unbounded system,

It is clear that $\kappa_1 \leq \tau_1 \leq \tau_2 \leq 2^{\aleph_0}$.

Theorem 2,1:

Let (Q, μ) be an \mathcal{L} -product of κ two-point spaces. The following properties are equivalent:

(1) (Q, μ) is a non- F -space.

(2) $\kappa \geq \tau_2$

Let (Q, μ) be a space, $A \subset Q$. We put $\mu^\circ A = A$, and for ordinal number α $\mu^\alpha A = \mu(\bigcup_{\beta < \alpha} \mu^\beta A)$.

Theorem 2,2:

Let an \mathcal{L} -product (Q, μ) of two-point spaces be a non- F -space and not countably compact. Then there exists $A \subset Q$ such that $\mu^\alpha A \neq \mu^{\alpha+1} A$ for all countable ordinal numbers α .

Definition:

We call a space (Q, μ) countably regular at a

point x , if x is an R -point $x)$ of every subspace P of (Q, u) such that $P = T \cup A \cup \{x\}$, $x \notin uT$, A is countable.

We call a space (Q, u) countably regular, if it is countably regular at each of its points.

Theorem 2,3^{xx}):

Let (Q, u) be an \mathcal{L} -product of κ two-point spaces, $\kappa \geq \tau_1$. Then (Q, u) and (Q, u^*) are not countably regular.

Theorem 2,4:

Let (Q, u) be an \mathcal{L} -product of κ two-point spaces. The following properties are equivalent:

- (1) For every $x \in Q$ there exists a closed set $T \subset Q$ such that $x \notin T$, $\text{card } T = \kappa_1$ and if U is a neighborhood of T in (Q, u) , then $x \in uU$.
- (2) (Q, u) is not regular.
- (3) (Q, u^*) is not regular.
- (4) $\kappa \geq \kappa_1$.

Problem: I do not know if τ_1 is the smallest cardinal number, satisfying the Theorem 2,3.

III. F -reduction.

Definition:

Let (Q, u) be a space, κ a cardinal number.

We denote by a symbol $\mathcal{F}_\kappa(Q, u)$ every collection $\{x_{\lambda, n}; \lambda \in \Lambda, n \in \mathbb{N}\}$ of elements of Q such that

- (1) $\text{card } \Lambda = \kappa$
- (2) There exists a point $x \in Q$ such that $x_{\lambda, n} \xrightarrow{n} x$ for every $\lambda \in \Lambda$.
- (3) If $\{m_i\} \in \mathbb{N}^\mathbb{N}$, $\{\lambda_i\} \in \Lambda^\mathbb{N}$, $\lambda_i \neq \lambda_j$ for $i \neq j$, then $x_{\lambda_i, m_i} \xrightarrow{i} x$.

x) x is an R -point of a space (Q, u) , if for every neighborhood U of x there exists its neighborhood V such that $uV \subset U$.

xx) cf [6], Theorem 1,1.

Theorem 3,1:

Let (Q, u) be an \mathcal{L} -product of n two-point spaces. Then (Q, u) is a discrete space if and only if there exists some $C_n(Q, u)$.

Theorem 3,2:

Let (Q, u) be an \mathcal{L} -product of n two-point spaces. Let $(P, v) = \mathcal{L}_{\alpha \in \Lambda} (P_\alpha, v_\alpha)$, and $\Lambda = n$, and $P_\alpha \subseteq n$ and every P_α contain at least two points.

If (Q, u) is a discrete space, (P, v) is also a discrete space.

Theorem 3,3:

Let (Q, u) be an \mathcal{L} -product of n two-point spaces. The following properties are equivalent:

(1) (Q, u) is discrete.

(2) $P_1 = T_1$

Theorem 3,4 x):

Let $(P, v) = \mathcal{L}_{\alpha \in \Lambda} (P_\alpha, v_\alpha)$, and $\Lambda = n^{P_0}$, and $P_\alpha \subseteq 2^{P_0}$, every P_α contain at least two points. Then (P, v) is a discrete space.

IV. The space of continuous functions.

Now we consider an \mathcal{L} -product (Q, u) and its subspaces, where $(Q, u) = \mathcal{L}_{\alpha \in \Lambda} (Q_\alpha, u_\alpha)$ and all (Q_α, u_α) are the spaces of real numbers E_1 (with a usual topology). We suppose that P is also a topological space and consider the space of real continuous functions on P .

In the following theorems $C(P)$ denotes the set of all real continuous functions on P , or the set of all real continuous and bounded functions on P , or the set of all mappings of P into $\langle 0, 1 \rangle$; $D(P)$ denotes any system of real functions on P . u denotes a topology on $C(P)$ (event. $D(P)$) such that $(C(P), u)$ (event. $(D(P), u)$) is a subspace of a given \mathcal{L} -product.

x) cf [6], Theorem 2,2.

Theorem 4,1:

Let \aleph be a cardinal number. Let $\text{card } f(P) \leq \aleph$ for every $f \in C(P)$. Then $\text{card } F(P) \leq \aleph$ for every continuous mapping f of P into any separable metric space.

This Theorem implies easily:

Theorem 4,2:

Let a set $f(P)$ be countable for every $f \in C(P)$. Then $(C(P), u)$ is an F -space.

Proposition II,3 in [6] implies easily the following Theorem:

Theorem 4,3:

Let $D(P)$ satisfy the following conditions:

- 1) If $g \in C(E_1)$, $f \in D(P)$, then $g \circ f \in D(P)$ ($g \circ f$ denotes the composition of f and g).
- 2) There exists a function $f \in D(P)$ such that $f(P)$ contains a closed subset which is dense-in-itself and non-meager. Then $(D(P), u)$ is a non- F -space.

This Theorem implies easily:

Theorem 4,4:

Let P be a compact space, containing an infinite discrete normally imbedded $x)$ subset. Then $(C(P), u)$ is a non- F -space.

V. F -reduction of a space of continuous functions.

Definition:

Let $D(P)$ be a system of real functions on P , $R(P)$ the system of all real functions on P , let \aleph be a cardinal number. The symbol $j_\aleph(D(P))$ denotes every collection $\{f_{\xi, m}; \xi \in \Xi, m \in N\}$ of elements of $D(P)$ such that

- 1) $\text{card } \Xi = \aleph$.
- 2) $f_{\xi, m} \xrightarrow{m} 1$ for all $\xi \in \Xi$.

x) A set R is said to be normally imbedded in a space P , if $R \subset P$ and every bounded continuous function on Q can be extended continuously to P . By discrete subset we mean simply a subset which, as a subspace, contains isolated points only.

3) if $\{a_n\} \in N''$, $\{b_n\} \in (C(P))^N$, $\{c_n\} \in \mathbb{R}^N$, $c_n \neq 0$,
for $i \neq j$, then $a_i \cdot b_{j,n} \rightarrow 1$.

Theorem 5,1:

Let $(C(P), \mu)$ be discrete, and $C(P) = \mathbb{R}$.
Then there exists $D_\mu(C(P))$. Let P contain a dense subset X , and $\text{card } X = \aleph_\alpha$, and let $j_\mu(C(P))$ exist there. Then $(C(P), \mu)$ is discrete.

Theorem 5,2:

Let $1 \in D(P)$. Let $D(P)$ satisfy

a) $f \in D(P) \Rightarrow \frac{f}{1+f} \in D(P)$

b) if $g \in C(E_1)$, g has all derivations, $g(0) = 0$, $g(1) = 1$, and $f \in D(P)$, then $g \circ f \in D(P)$.

Then the following propositions are equivalent:

(1) there exists $D_{\mu_0}(D(P))$.

(2) there exists $j_{\mu_0}(D(P))$.

Theorems 5,1 and 5,2 imply easily the Theorem II, 1 in [6].

Theorem 5,3:

Let P be a space, containing a dense countable metrizable subset. Let every neighborhood of every point $x \in P$ contain a neighborhood of x , which is a dense-in-itself non-meager normal space.

Let $D(P) \subset C(P)$ such that:

1) $\mu D(P) = C(P)$ (i.e.: for every $f \in C(P)$ there exist $f_n \in D(P)$, $n = 1, 2, \dots$ such that $f_n \xrightarrow{\mu} f$).

2) if $A \subset P$ is closed, $y \notin A$, then there is a function $f \in D(P)$ with $f(y) = 0$, $f(x) = 1$ for all $x \in A$.

3) if $f, g \in D(P)$, then $f \cdot g \in D(P)$.

Then for any $f \in C(P) - (0)$ there exists a set $H_f \subset D(P)$ such that $\mu H_f = C(P) - (f)^x$.

If more

4) there exists a function $h \in C(P)$ such that $h \neq 0$, and $h - f \in D(P)$ for all $f \in D(P)$,

x) The closure μH_f of H_f we certainly consider in the space $(C(P), \mu)$ only. Some non-continuous functions are the limits of sequences of points of H_f , too.

then there exists $H_0 \in D(P)$ such that
 $\mu H_0 = C(P) - (0)$.

This Theorem may be applied, for example, for the set of all real functions of real variables, having all derivations.

VI. Some results about the space $(C(P), \mu)$.

Theorem 6,1:

If a normal space P contains a locally finite disjoint system, the power of which is \aleph_1 (event. \aleph_1 , event. $\aleph_1 \cdot \aleph_2$), then $(C(P), \mu)$ and $(C(P), \mu^*)$ are not regular (event. $(C(P), \mu)$ and $(C(P), \mu^*)$ are not countably regular, event. $(C(P), \mu)$ contains a set H such that $\mu^\alpha H \neq \mu^{\alpha+1} H$ for all countable ordinal numbers α).

Theorem 6,2:

Let ω be the smallest ordinal number, the power of which is a regular cardinal number \aleph . Let every subspace of some space (Q, μ) contain a dense subset, the power of which is $< \aleph$.

Then there exists an ordinal number α for every $R \in Q$ such that $\alpha < \omega$ and $\mu^\alpha R = \mu^{\alpha+1} R$.

Theorem 6,3:

Let P be a union of the countable number of compact metric spaces. Then every subspace of $(C(P), \mu)$ contains a dense countable subset.

Theorem 6,4 which is a strengthening of Theorem 1,2 in [2], follows immediately from the Theorems 6,2 and 6,3:

Theorem 6,4:

Let P be a union of the countable number of compact metric spaces.

Then, for every $H \in C(P)$, $\mu^\alpha H = \mu^{\alpha+1} H$ for some countable α and consisting from open sets, \forall

VII. Countable compactness of $(C(P), \mu)$.

$C(P)$ denotes the set of all continuous mappings of P into $\langle 0, 1 \rangle$ in this section.

Theorem 7,1:

Let a space P contain a normally imbedded discrete set of the power \bar{c} . Then $(C(P), u)$ is not countably compact.

Theorem 7,2:

Let a normal space P contain a closed G_δ subset which is not open. Then $(C(P), u)$ is not countably compact.

Theorem 7,3:

Let $\bar{c} = \aleph_1$.

- a) If P is a perfectly normal space, then $(C(P), u)$ is countably compact only for a countable discrete P .
- b) If P is a normal space and $(C(P), u)$ is a non- F -space, then $(C(P), u)$ is not countably compact.

VIII. Borel functions.

Let P be a perfectly normal space. $B(P)$ denotes the set of all real Borel functions on P , or the set of all real bounded Borel functions (or bounded by a certain constant), or the set of all characteristic functions of Borel subsets of P . A definition of the topology u on $B(P)$ is evident.

Theorem 8,1:

Let us suppose that a perfectly normal space P contains a normally imbedded discrete subset, the power of which is $\bar{c} = \aleph_1$, let $\text{card } P \leq 2^{\aleph_1}$. Then $(B(P), u)$ is a discrete space.

Theorem 8,2:

If a perfectly normal space P contains a Borel subset, which may be mapped continuously on a topological product of \aleph_1 two-point spaces, and $\text{card } P \leq 2^{\aleph_1}$, then $(B(P), u)$ is a discrete space.

It is clear that the Theorems 6,1; 6,3; 6,4; 7,1 hold also for $(B(P), u)$.

The problem, raised by J. Novák, whether $(B(E_\gamma), u)$ is regular, remains unsolved. It may be shown only, that there

exists a subspace P of E_1 such that $(B(P), u)$, and $(B(P), u^*)$ are not regular (neither are they countably regular).

R e f e r e n c e s.

- [1] E. Čech: Topologické prostory, Praha 1959.
- [2] M. Neubauer: Sur l'espace des fonctions continues, Fund.Math. 31 (1938), 269 - 273.
- [3] J. Novák: Sur les espaces (\mathcal{L}) et sur les produits cartesiens (\mathcal{L}) , Spisy vydávané přírodověd. fakultou MU, Brno 1939.
- [4] J. Novák, L. Mišík: O \mathcal{L} -priestorach spojitých funkcií. Mat.fyz.sborník SAVU 1 (1951), 1 - 13.
- [5] B. Pospíšil: Sur les fonctions continues, Fund. Math. 31 (1938), 262 - 268.
- [6] V. Šedivá: On pointwise convergence of sequences of continuous functions, CMUC 1,2 (1960), 43 - 51.